

Table 1. Estimated numbers of cases of HIV/AIDS, by year of diagnosis and selected characteristics of persons, 2001–2004—35 areas with confidential name-based HIV infection reporting

	Year of diagnosis			
	2001	2002	2003	2004
Age at diagnosis (yrs)				
< 13	360	288	203	174
13–14	53	46	53	41
15–19	1,012	1,023	1,027	1,080
20–24	3,241	3,154	3,345	3,762
25–29	4,608	4,499	4,290	4,604
30–34	6,776	6,305	5,688	5,527
35–39	8,261	7,719	7,112	6,754
40–44	6,699	6,395	6,483	6,344
45–49	4,709	4,486	4,514	4,536
50–54	2,735	2,643	2,733	2,776
55–59	1,411	1,330	1,414	1,594
60–64	730	763	705	804
≥65	674	627	623	734
Race/ethnicity				
White, not Hispanic	11,242	11,352	11,097	11,806
Black, not Hispanic	21,556	20,237	19,310	19,206
Hispanic	7,714	6,964	7,078	6,970
Asian/Pacific Islander	279	319	367	394
American Indian/Alaska Native	171	202	187	208
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	16,625	16,852	16,804	18,203
Injection drug use	5,171	4,379	4,177	3,828
Male-to-male sexual contact and injection drug use	1,525	1,431	1,398	1,372
Heterosexual contact	5,095	4,843	4,720	4,581
Other ^a	214	183	179	161
Subtotal	28,630	27,689	27,279	28,143
Female adult or adolescent				
Injection drug use	2,877	2,408	2,252	2,134
Heterosexual contact	9,192	8,709	8,248	8,102
Other ^a	211	187	205	174
Subtotal	12,280	11,303	10,706	10,410
Child (<13 yrs at diagnosis)				
Perinatal	306	245	186	145
Other ^b	54	44	18	32
Subtotal	360	288	204	177
Subtotal for 33 states with confidential name-based HIV infection reporting	41,207	39,222	38,139	38,685
Subtotal for U.S. dependencies, possessions, and associated nations with confidential name-based HIV infection reporting	63	58	49	45
Total^c	41,270	39,280	38,188	38,730

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

Data include persons with a diagnosis of HIV infection. This includes persons with a diagnosis of HIV infection only, a diagnosis of HIV infection and a later AIDS diagnosis, and concurrent diagnoses of HIV infection and AIDS.

Since 2000, the following 35 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, Guam and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.

^a Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^b Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^c Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.